

A211 029

Are ozone exceedance rates decreasing? Comment on "Extreme value analysis of environmental time-series: An application to trend detection in ground-level ozone" by R.L. Smith

Adrian E. Raftery  
University of Washington

ABSTRACT

R.L. Smith (1989) in his *Statistical Science* discussion paper, proposed new methods for analyzing extreme values based on the point process view of high-level exceedances, and illustrated them with a detailed analysis of ozone data from Houston, Texas. The methods are powerful and, in particular, the point process of cluster peaks over a high threshold provides a remarkable condensation of the massive data set that he analyzes. It involves little loss of relevant information and permits fairly simple analyses.

Smith's conclusion is that there is no trend in the overall levels of the series, but that there is a marked downward trend in the extreme values. It seems hard to find physical explanations for this, and here the evidence is reassessed in terms of a comparison between competing models for the intensity of a Poisson process. This suggests that there is some evidence for a decreasing trend in exceedance rates but that it is rather weak. If there is a trend, it seems more likely to consist of a fairly abrupt change than a gradual decrease. The possibility that such a change is due to an improvement in measurement technology is discussed. The possibility of long-memory dependence is also considered and the clustering method used is discussed.

DTIC  
ELECTE  
AUG 09 1989  
S B D

Adrian E. Raftery is Associate Professor, Department of Statistics, GN-22, University of Washington, Seattle, WA 98195. This research was supported by the Office of Naval Research under Contract no. N-00014-88-k-0265. The author is grateful to Richard L. Smith for providing the ozone data on which the analyses are based, to Peter Guttorp for very helpful discussions, and to Michael A. Newton for useful comments. The computer programs used to carry out the analyses may be obtained from the author by sending electronic mail to [raftery@entropy.ms.washington.edu](mailto:raftery@entropy.ms.washington.edu).

DISTRIBUTION STATEMENT A  
Approved for public release;  
Distribution Unlimited

## 1. INTRODUCTION

In his excellent paper, Smith (1989) has synthesized a range of powerful methods for the analysis of extreme values. The point process of cluster peaks over a high threshold provides a remarkable condensation of the massive data set that he analyzes. It involves little loss of relevant information and permits fairly simple analyses. The methodology is sure to find wide application.

Nevertheless, I find it hard to think of physical explanations for the conclusion that there has been a downward trend in the extreme values without any accompanying decrease in the overall levels of the ozone series. Here I try to reassess the evidence in terms of a comparison between competing models for the intensity of a Poisson process. The analysis suggests that there is some evidence for a decreasing trend in exceedance rates but that it is rather weak. If there is a trend, it seems more likely to consist of a fairly abrupt change than a gradual decrease. The possibility that such a change is due to an improvement in measurement technology is discussed. I also consider the possibility of long-memory dependence and discuss the clustering method used.

## 2. ARE OZONE EXCEEDANCE RATES DECREASING?

The evidence in Smith (1989) for decreasing exceedance rates consists mainly of the fact that the estimated trend was downward in all the models that incorporated a trend. However, these models did not appear to fit better than models that did not incorporate a trend. For example, the likelihood ratio test statistic for splitting the data was 16.6 with 18 degrees of freedom.

This may be due more to the large number of degrees of freedom than to the absence of an effect. It might be worth, for example, fitting a model of the form (4.1), but with  $\mu_{ij} = \alpha_j + \beta\delta_i$ , where  $\delta_i = 0$  for 1973-80 and  $\delta_i = 1$  for 1981-86. One could then test the hypothesis that  $\beta = 0$ , which involves only one degree of freedom rather than 18. There are many other parsimonious possibilities.



<input checked="checked" type="checkbox"/>	
<input type="checkbox"/>	
<input type="checkbox"/>	
Codes	
Dist	Avail and/or Special
A-1	

*Non-homogeneous Poisson process models for exceedances*

Professor Smith's conclusion corresponds to a decreasing rate of occurrence in the point processes of exceedances above high thresholds. This process was not fully observed, and the proportion of time monitored varied over the period, increasing gradually but significantly. I therefore expressed times of occurrence in terms of monitored time since the start of the data, rather than calendar time. Also, ozone levels are highly seasonal. I estimated the seasonal effect as piecewise constant within each of the six 61-day periods, and deseasonalized the data by transforming the time axis (Cox and Lewis, 1966). The resulting series of events are shown in Tables 1 and 2. I denote by  $T$  the period of observation and by  $t = (t_1, \dots, t_n)$  the event times.

TABLE 1  
Times of occurrence of exceedances above  
threshold level 16, in monitored days from  
the start of the data, deseasonalized  
(read down the columns)

62	511	913	1442	1760	2168	2890	3550
71	539	934	1484	1774	2204	2976	3565
88	555	1006	1506	1830	2245	3051	3636
114	585	1022	1527	1893	2363	3079	3685
122	596	1043	1554	1924	2417	3153	3827
146	610	1057	1585	1986	2465	3257	3846
184	627	1138	1617	2007	2535	3302	3891
219	662	1184	1621	2045	2568	3325	
235	740	1231	1624	2076	2608	3346	
285	779	1318	1642	2099	2705	3403	
327	847	1409	1742	2122	2791	3421	

If there is no trend, the data in Tables 1 and 2 are very nearly from a homogeneous Poisson process; we denote this model by  $M_0$ . This assumes that any short-term correlation has been removed by considering only cluster peaks. An alternative hypothesis is that the exceedance rate has been decreasing smoothly and gradually. This may conveniently be represented by the log-linear Poisson process,  $M_1: \lambda(s) = \rho e^{-\beta s}$ , where  $\lambda(s)$  is the rate of occurrence at time  $s$ . Another possibility, suggested by the splitting of the data in Smith (1989), is that the exceedance

TABLE 2  
Times of occurrence of exceedances above  
threshold level 20, in monitored days from  
the start of the data, deseasonalized  
(read down the columns)

71	327	1057	1624	1986	2122	2705	3079
88	555	1442	1642	2045	2417	2791	3550
122	847	1484	1830	2076	2465	2890	3846
285	913	1585	1893	2099	2608	2976	

rate decreased fairly abruptly within a short time period. This may be represented by the change-point Poisson process,  $M_2$ :  $\lambda(s) = \lambda_1$  if  $0 \leq s \leq \tau$  and  $\lambda(s) = \lambda_2$  if  $\tau < s \leq T$ .

#### Model comparison

The three competing models,  $M_0$ ,  $M_1$  and  $M_2$ , may be compared using the Bayes factor, or ratio of posterior to prior odds for  $M_i$  against  $M_j$ ,  $B_{ij}$ , for each pairwise comparison. It has been argued that Bayes factors are better measures of evidence than P values (Berger and Sellke, 1987), and they are also more readily applicable to the comparison of non-nested models.

With vague prior information, the Bayes factor for  $M_0$  against the log-linear Poisson process,  $M_1$ , is

$$B_{01} = 0.645(n-1) / \int_0^\infty e^{-Ry} \left[ \frac{y}{1-e^{-y}} \right]^{n-1} dy, \quad (1)$$

where  $R = \sum t_i / T$  (Akman and Raftery, 1986a). The Bayes factor against the change-point Poisson process,  $M_2$ , is

$$B_{02} = 4\sqrt{\pi} \Gamma(n+1/2) / \sum_{i=0}^n \Gamma(i+1/2) \Gamma(n-i+1/2) I_i \quad (2)$$

(Raftery and Akman, 1986). In (2),  $I_i = \int_{u_i}^{u_{i+1}} x^{-(i+1/2)} (1-x)^{-(n-i+1/2)} dx$ , where  $u_i = t_i / T$ .

A classical test statistic for  $M_0$  against  $M_1$  is  $U = (R - 1/2n) / \sqrt{\frac{n}{12}}$ , which has approximately a standard normal distribution under  $M_0$  (Cox and Lewis, 1966). A test of  $M_0$  against  $M_2$  may be based on the quantity

$$\Delta = n^{-1/2} \max_{.01 \leq u_i \leq .99} \{ |g(i-1, u_i)|, |g(i, u_i)| \}, \quad (3)$$

where  $g(i, u) = i \sqrt{\frac{1-u}{u}} - (n-i) \sqrt{\frac{u}{1-u}}$ . The approximate 5% critical value for this test is 3.29 (Akman and Raftery, 1986b).

In Table 3 there is evidence against a gradual decrease in exceedance rate of the form specified by  $M_1$ . The posterior odds for the change-point Poisson process are 2.7:1 and 2:1 at threshold levels of 16 and 20 respectively. In the words of Jeffreys (1961), this constitutes evidence against the homogeneous Poisson process, but it is not worth more than a bare mention. In addition, the result of the classical test based on (3) is not significant.

TABLE 3  
Model comparison results.  
 $B_{01}$  is the Bayes factor for the homogeneous Poisson process  
against the log-linear Poisson process given by (1);  
 $U$  is the corresponding classical test statistic.  
 $B_{02}$  is the Bayes factor for the homogeneous Poisson process  
against the change-point Poisson process given by (2);  
 $\Delta$  is a corresponding classical test statistic given by (3).

	Threshold level	
	16	20
$\log_{10} B_{01}$	1.69	1.11
$U$	-1.02	-0.97
$\log_{10} B_{02}$	-0.44	-0.30
$\Delta$	2.33	2.15

### *Checking the homogeneous Poisson process*

Since the evidence for a trend appears weak, it seems worth checking the homogeneous Poisson process model itself. One way of doing this is to compare the observed evolution with those of several data sets simulated from the model. Since the homogeneous Poisson process is time-reversible, we can do the same with the time-reversed data set. Ripley (1977), who pioneered this approach, used point estimates of the model parameters in the simulations, but this may lead to simulated bands which are too narrow. Here, uncertainty about the parameter  $\lambda$  of the homogeneous Poisson process is incorporated as follows (Rubin, 1984; Raftery, 1988). First, generate a value of  $\lambda$  from its posterior distribution, taken here to be Gamma  $(n+1/2, T)$ , and then proceed as before.

The result is shown in Figure 1. The data do not go outside the simulated bands except very briefly in Figures 1(b) and 1(d). Once again, the evidence against the homogeneous Poisson process does not seem strong. The simulated bands may still be too narrow because they do not take account of uncertainty about the estimated seasonal effect, and so the evidence may be even weaker than it appears.

### *Analysis of the change-point Poisson process model*

The analysis so far suggests that, if there is a trend, it is better represented by a change-point than by a gradual decrease. The posterior distribution of the change-point is given by equation (2.3) of Raftery and Akman (1986) and is shown in Figure 2. At threshold level 20, the posterior distribution is less diffuse than at level 16. The posterior mode is April 11, 1984, which is at the beginning of the 1984 "ozone season". This result reflects the fact that there was only one exceedance above level 20 in each of 1984, 1985 and 1986, compared with 28 such exceedances in the previous ten seasons. The posterior mode at threshold level 16, which is less marked than at level 20, is June 25, 1981.

The analysis here is tentative in many ways. In particular it seems important to include relevant covariates, especially temperature, as emphasized by Davison and Hempill (1986).






FIGURE 1. Diagnostic checking for the homogeneous Poisson process. In each graph, the solid line represents the data and the dotted lines are the outer envelopes of 19 simulations from the model, as described in the text.



(a) Threshold level 16



(b) Threshold level 20

FIGURE 2. Posterior distribution of the change-point in the change-point Poisson process model.



It would be interesting to know if there were any events which could have caused an abrupt change, such as legislation, changes in Federal standards, highway development, or changes in data collection, checking and reporting practices.

My colleague Peter Guttorp has suggested the following as a possible explanation. In the past, hourly ozone measurements were often based on 25-minute averages. Now it is more usual for measurement to be continuous, so that hourly measurements are based on 60-minute averages. Such a change in instrumentation would not have changed the overall level of the series, but it might well have reduced its variability, and hence the exceedance rate, because the measurements are based on more data. This would have been an abrupt change rather than a gradual one, and so seems consistent with the results here and in Smith (1989). It would be interesting to know if there was such a change in instrumentation in Houston during the period covered by the data. If so, it would suggest that we are not seeing an improvement in compliance with the Federal standard, but rather a change in measurement technology.

## 2. OTHER ISSUES

### *Long-memory dependence*

Long-memory dependence is known to be a feature of at least some climatic variables (Haslett and Raftery, 1989, and references therein). Climate influences ozone levels, so it is possible that ozone levels may also exhibit long-memory dependence. This is characterized by small but non-negligible autocorrelations at long lags, an infinite spike at zero in the spectrum or "cycles of all periods", and high variability of the sample mean and other statistics. It is hard to detect but can dramatically affect statistical analyses.

Is there any evidence of long-memory dependence in the ozone data? How could it be detected? Would it affect the analysis of extreme values? If so, how could it be incorporated into the analysis?

### *The clustering method*

The method for forming clusters used by Smith (1989) is essentially the single link method. This has the possible disadvantage that two clusters six days apart with a single exceedance between them could be merged (Gordon, 1981). An agglomerative sum of squares method might be preferable given that the aim is to obtain compact clusters. Inspection of the dendrogram could help with the choice of a cluster interval.

### REFERENCES

- Akman, V.E. and Raftery, A.E. (1986a). Bayes factors for non-homogeneous Poisson processes with vague prior information. *J. R. Statist. Soc., ser. B* **48** 322-329.
- Akman, V.E. and Raftery, A.E. (1986b). Asymptotic inference for a change-point Poisson process. *Ann. Statist.* **14** 1583-1590.
- Berger, J.O. and Sellke, T. (1987). Testing a point null hypothesis: the irreconcilability of P values and evidence. *J. Amer. Statist. Ass.* **82** 112-122.
- Cox, D.R. and Lewis, P.A.W. (1966). *The Statistical Analysis of Series of Events*. Methuen, London.
- Davison, A.C. and Hempill, M.W. (1986). On the statistical analysis of ambient ozone data when measurements are missing. *Atmospheric Environment* **21** 629-639.
- Gordon, A.D. (1981). *Classification*. Chapman and Hall, London.
- Haslett, J. and Raftery, A.E. (1989). Space-time modelling with long-memory dependence: Assessing Ireland's wind power resource (with Discussion). *Appl. Statist.* **38** 1-50.
- Jeffreys, H. (1961). *Theory of Probability* (3rd ed.). Clarendon Press, Oxford.
- Raftery, A.E. (1988). Analysis of a simple debugging model. *Appl. Statist.* **37** 12-22.
- Raftery, A.E. and Akman, V.E. (1986). Bayesian analysis of a Poisson process with a change-point. *Biometrika* **73** 85-89.

Ripley, B.D. (1977). Modelling spatial patterns (with Discussion). *J. R. Statist. Soc., ser. B* **39** 172-212.

Rubin, D.B. (1984). Bayesianly justifiable and relevant frequency calculations for the applied statistician. *Ann. Statist.* **12** 1151-1172.

Smith, R.L. (1989). Extreme value analysis of environmental time series: An application to trend detection in ground-level ozone. *Statist. Sci.* **4**, to appear.

REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM
1. REPORT NUMBER 165	2. GOVT ACCESSION NO.	3. RECIPIENT'S CATALOG NUMBER
4. TITLE (and Subtitle) Are ozone exceedance rates decreasing? Comment on "Extreme Value Analysis of Environmental Time-Series: An application to trend detection in ground-level ozone" by R.L. Smith		5. TYPE OF REPORT & PERIOD COVERED Technical Report 1/1/89-7/31/89
7. AUTHOR(s) Adrian E. Raftery		6. PERFORMING ORG. REPORT NUMBER
9. PERFORMING ORGANIZATION NAME AND ADDRESS Department of Statistics, GN-22 University of Washington Seattle, WA 98195		8. CONTRACT OR GRANT NUMBER(s) N00014-88-K-0265 P00001
11. CONTROLLING OFFICE NAME AND ADDRESS ONR Code N63374 1107 NE 45th Street Seattle, WA 98105		10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS NR-661-003
14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office)		12. REPORT DATE July 1989
		13. NUMBER OF PAGES 11
		15. SECURITY CLASS. (of this report) Unclassified
		15a. DECLASSIFICATION/DOWNGRADING SCHEDULE
16. DISTRIBUTION STATEMENT (of this Report)  APPROVED FOR PUBLIC RELEASE: DISTRIBUTION UNLIMITED.		
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report)		
18. SUPPLEMENTARY NOTES		
19. KEY WORDS (Continue on reverse side if necessary and identify by block number) Non-homogeneous Poisson process; Seasonal effect; Bayes factor; Non-nested model; Change-point; Log-linear Poisson process; Bayesian model checking; Long-memory dependence; Clustering algorithms.		
20. ABSTRACT (Continue on reverse side if necessary and identify by block number) R.L. Smith (1989) in his Statistical Science discussion paper, proposed new methods for analyzing extreme values based on the point process view of high level exceedances, and illustrated them with a detailed analysis of ozone data from Houston, Texas. The methods are powerful and, in particular, the point process of cluster peaks over a high threshold provides a remarkable condensation of the massive data set that he analyzes. It involves little loss of relevant information and permits fairly simple analyses.		

Smith's conclusion is that there is no trend in the overall levels of the series, but that there is a marked downward trend in the extreme values. It seems hard to find physical explanations for this, and here the evidence is reassessed in terms of a comparison between competing models for the intensity of a Poisson process. This suggests that there is some evidence for a decreasing trend in exceedance rates but that it is rather weak. If there is a trend, it seems more likely to consist of a fairly abrupt change than a gradual decrease. The possibility that such a change is due to an improvement in measurement technology is discussed. The possibility of long-memory dependence is also considered and the clustering method used is discussed.